

Stephen Kriss MPhil FCPodS

Consultant Podiatric Surgeon

ESSENTIAL PREOPERATIVE INFORMATION FOR PATIENTS
FACULTY OF PODIATRIC SURGERY

This is a generic document for all patients undergoing foot surgery. There is a lot of information in this guide some of which may not be applicable to the surgery you are having. The purpose of this document is to inform you. You will receive other literature specifically tailored to the surgery you are having. If you have any queries please contact me to discuss

THIS INFORMATION IS INTENDED AS A GUIDE ONLY

It is designed to be considered in conjunction with specific information concerning your particular operation

Instructions to patients

Please take time to read this booklet well before your planned foot operation. It is your responsibility to ask, if you feel you need more information about your foot condition or any proposed treatment.

1 What is the purpose of this booklet?

This booklet has been designed by a working party of Podiatric Surgeons. It is intended to provide you with additional information about your proposed treatment and assist you in making an informed choice. Information specific to your own planned operation will be provided separately.

2 Who is responsible for my care?

Your treatment / surgery will be undertaken / supervised by a Podiatric Surgeon who is a Fellow of the Faculty of Surgery (College of Podiatrists). Podiatric Surgeons are not registered medical practitioners (Medical Doctors), but are non-medical specialists in the surgical and non-surgical management of problems of the foot and associated structures.

Titles:

The podiatric surgery team may consist of several different professionals all dedicated to the success of your surgery and for your assistance the common titles are explained below.

Podiatrist:

Is a clinician who has studied for three years to obtain a degree in podiatric medicine and registration with the Health Professions Council. Podiatrists are independent clinicians, qualified to diagnose and treat foot problems. Podiatrists may specialise in particular areas of work e.g. the care of the diabetic patient or sports medicine. With the exception of nail surgery, podiatrists undertake the treatment of foot problems by non-invasive methods (until recently podiatrists were known as chiropodists).

Trainee in Podiatric Surgery:

Is a podiatrist who has studied for a further two or three years to complete the initial sections of the Faculty of Surgery exams (or obtained an MSc degree) in the theory of podiatric surgery and is undertaking a formal surgical training programme under the supervision of a Consultant Podiatric Surgeon who is an accredited Tutor of the Faculty of Surgery. The aim of the trainee is to successfully pass the Fellowship exam and continue through their post Fellowship period.

Specialist Registrar in Podiatric Surgery:

Is a past trainee in podiatric surgery who has successfully completed the gained his / her Podiatric Surgical Fellowship (qualification in the practice of podiatric surgery) and is working as part of a continued training programme towards completion of the three year post Fellowship specialist training period.

Podiatric Surgeon:

A Podiatric Surgeon has successfully gained both his / her Fellowship and a certificate of completion of training. A podiatric surgeon is a non-medically qualified specialist in the treatment of foot problems by both surgical and non-surgical methods. A podiatric surgeon has completed the training process and may have his / her own caseload. .

Consultant Podiatric Surgeon:

After some years of practice within a Health Service Department of Podiatric Surgery, a podiatric surgeon may be appointed as a consultant i.e. the lead clinician appointed by an NHS Trust to provide a podiatric surgery service.

THIS INFORMATION IS INTENDED AS A GUIDE ONLY

It is designed to be considered in conjunction with specific information concerning your particular operation

3 Do I have a choice with regard to my anaesthetic for foot surgery?

Foot surgery can be undertaken using a number of types of anaesthetic, these are:

- ✓ Local anaesthetic
- ✓ Local anaesthetic + sedation
- ✓ Spinal anaesthetic
- ✓ General anaesthetic

Not all of these types of anaesthetic may be suitable for you and not all are offered in all centres. You should discuss your preferences with one of the team during your visits to the department.

4 *Local Anaesthetic*

Local anaesthetic means you can be awake during your operation though your foot will be numb and the operation will be screened away from you. You will be able to listen to music or read whilst your operation is done. Many patients choose to have their foot operation using a local anaesthetic. Sometimes it is necessary to give you additional local anaesthetic during your surgery (about 4 cases per 100).

Local Anaesthetic with Sedation

Sedation normally requires an anaesthetist to be present to administer a sedative through a small needle placed in a vein in the back of the hand. Sedation usually has the benefit of reducing anxiety during the operation. You will not be able to eat or drink for approximately 6-8 hours prior to your surgery.

General Anaesthetic

With a general anaesthetic you are put to sleep though you will still require a local anaesthetic (this is usually given once you are asleep). Some patients can feel sick when they wake-up. You will not be able to eat or drink for approximately 6-8 hours prior to your surgery.

5 Can I eat and drink as usual before my operation?

If your surgery is going to be undertaken using a local anaesthetic only, then you are able to eat and drink as normal. If your surgery is planned under general anaesthetic / sedation then you should not eat or drink for six hours prior to your operation. If you have any questions about this, please speak to one of the Podiatric Surgery team.

6 What will I wear on my foot after my operation?

Where required we provide patients with a special post-operative shoe which is designed to accommodate the bandages over the foot,

THIS INFORMATION IS INTENDED AS A GUIDE ONLY

It is designed to be considered in conjunction with specific information concerning your particular operation

7 Will I be in a cast after surgery or have to use crutches?

This will depend on the type of operation you have had. You will be told beforehand if crutches or a cast are required following your operation. You will be shown how to safely use crutches if these are necessary.

8 What do I need to do before I come in for my operation?

- Check your appointment letter to make sure you know what time to arrive and where to go
- Cut your toenails and clean under the nails well the day before surgery.
- Remove all nail polish.
- Remove jewellery except for your wedding ring (if you have one)
- Wear loose clothing as a large dressing will be applied after the operation.
- You may bring along a personal stereo player with headphones
- Please make sure you arrange transport to and from the hospital / day-surgery unit. You must have a fit and responsible adult to stay with for at least the first night of your operation.
- You should arrange the recommended period of rest for after your operation as advised by your podiatric surgeon
- If you are under 16yrs of age you must have a parent or legal guardian accompany you to all your appointments
- If you will need to take your regular pre-prescribed medication during your time in the department, or may have need of an inhaler or angina spray or EpiPen; please remember to bring these with you.
- If you have been advised to arrange any tablets / medicines by your Podiatric Surgeon remember to obtain these **prior** to your operation. If you have been asked to obtain these from your own doctor please arrange this well in advance.
- Take any prescribed medicines as normal unless you have been told differently.
- Read any additional information you may be given regarding your admission for surgery.

9 What will happen on the day of my operation?

- You will be asked to come in to the hospital / day-surgery unit on the day of your operation;
- When you arrive you will be welcomed by one of the team who will escort you to the admission area where you may be asked to change into a theatre gown;
- Your consent form will be checked by one of the Podiatric Surgery team and the proposed operation will be fully re-explained. This is an opportunity for you to ask further questions;
- You may be asked more questions about your health & have your blood pressure checked,
- The operation site will be clearly marked;
- If you are going to be provided with crutches after your surgery, you will be shown how to use these either before or after your operation;
- If you are having a local anaesthetic this will be given before you are taken to theatre. If you are having general anaesthetic, the anaesthetist will come and see you before your operation;
- Your local anaesthetic will be checked for effectiveness by one of the team before you are taken into theatre;
- When the team is ready, you will be conveyed into the operating theatre for your surgery;
- Your foot will be cleaned with surgical scrub and sterile sheets will be placed around the area;
- Your local anaesthetic will be re-checked to make sure the area of your foot is fully numb
- A tourniquet (like a blood pressure cuff around your ankle) may be used during your operation. This may be uncomfortable for a short while until you ankle tissues adjust, but most people find it quite acceptable after about 5 minutes;
- After your operation your foot will be bandaged up carefully and you will be transferred back to a recovery area.

THIS INFORMATION IS INTENDED AS A GUIDE ONLY

It is designed to be considered in conjunction with specific information concerning your particular operation

10 What will happen after I have my operation?

- You will be transferred from theatre to a recovery area;
- One of the team will make sure that you are comfortable and check your dressing
- It is usual for you to be monitored for about 15-30 minutes after your operation, this varies depending on what operation(s) you have had done and the type of anaesthetic. You may be offered a drink and something to eat.
- A special post-operative shoe or boot (or sometimes a cast) will be applied to the foot;
- One of the team will check you are pain-free, read through and explain your postoperative instructions, including any contact numbers to use in case of a problem;
- You will be discharged home with your escort (who must be a responsible adult).

11 Can I go home the same day?

Most foot surgery can be done as a “day case operation” which means that are able to go home the same day. You will need a fit and responsible adult to stay with you for at least the first night after your operation. The amount of help patients need after surgery varies a great deal. You must make sure you fully understand the impact of your proposed surgery by discussing this with one of the Podiatric Surgery team.

12 What happens if I have a problem after my operation?

You will be provided with an advice sheet to take home after your operation detailing your postoperative instructions, and useful information including any contact numbers to use in case of a problem.

13 Will I need to take time off work after my surgery?

You are advised to rest after surgery. The amount of time required varies according to the type of operation and your own circumstances. Please refer to the specific operation information leaflet provided by your podiatric surgeon or ask one of the clinic staff.

14 Will I have to come back to hospital / clinic after my surgery?

The arrangements for follow-up after Podiatric Surgery will vary from service to service but you will be informed of what to expect before your operation. In most cases you will be reviewed by one the Podiatric Surgery team following your operation though your initial postoperative care may be via your own GP surgery. .

THIS INFORMATION IS INTENDED AS A GUIDE ONLY

It is designed to be considered in conjunction with specific information concerning your

particular operation
Page 5 of 11

THINGS TO CONSIDER BEFORE HAVING A FOOT OPERATION

Patients seldom have problems following podiatric surgery and most are pleased with their result. Unfortunately all treatments/operations carry risks. You need to carefully consider the severity of your problem against the likely risk/benefit from any proposed surgery (for more information on understanding risks, see the BestTreatments website (<http://www.besttreatments.co.uk/btuk/howtouse/7432.html>)). This information booklet is not intended to worry you. It is hoped that by providing as much information as possible, all your pre-surgery questions are answered. If you have additional questions do not hesitate to ask a member of the team. For convenience this section is laid out in alphabetical order.

Activity

You will be required to rest after your operation. How much rest and how much you are allowed to undertake will be discussed with you before and after your surgery. Postoperative recovery times vary between patients. For an indication of this please refer to the specific operation information leaflet provided by your podiatric surgeon or ask one of the clinic staff.

Alcohol Consumption

You are advised to avoid alcohol after your foot surgery, whilst on medication. Alcohol may interact with one or more of your medicines in addition to increasing the risk of falls postoperatively.

Altered Walking Pattern

Foot surgery may affect the way you walk. Following foot surgery, patients naturally favour the other foot. This is usually temporary, until the foot settles and you become accustomed to walking normally again. Sometimes we provide a foot splint to help with symptoms.

Avascular Necrosis

This is where part of the bone loses its blood supply. This may occur after an operation or even from simple trauma. The bone may ‘weaken’ and change shape. Usually the blood supply will return to the bone with time. Sometimes the bone is damaged resulting in problems such as secondary arthritis. This is a very rare occurrence.

Being awake during the operation

If you have elected to have your operation under local anaesthetic you will be awake during your operation. Your foot / leg will be tested before you enter theatre to ensure you are pain free. We check this again once you are in theatre. In the rare event you feel any discomfort you simply need to notify the nurse and we will be able to deal with this immediately. This is a rare event and most patients are able to chat to one of the nursing team or listen to their music whilst the operation is being performed.

Bleeding

There will be some bleeding postoperatively but the majority of this is controlled during the operation. Infrequently, a small collection of blood in the deeper tissues may form. This occurs less than 1% of the time and can be dealt with and generally settles without problems.

Changing Your Mind

Whilst we will have done our very best to diagnose your foot complaint and formulate an appropriate treatment plan for you, you do **not** have to follow this treatment plan against your wishes. Patients have the right to change their mind at any time about their treatment. Please be assured that should you change your mind about your proposed treatment staff will do their utmost to find an alternative treatment plan that suits you better.

THIS INFORMATION IS INTENDED AS A GUIDE ONLY

It is designed to be considered in conjunction with specific information concerning your particular operation

Chronic Regional Pain Syndrome (Reflex Sympathetic Dystrophy - RSD)

This is a rare condition, which may occur even after even minor injury. It is not always associated with surgery. The condition is very uncommon and can occur after any type of operation. Persistent / chronic pain develops as well as swelling / colour changes and changes to bones of the affected limb. Treatment requires early diagnosis and referral to a pain specialist. Complete resolution of RSD may not always be possible. This is not a predictable event but fortunately uncommon (0.1%).

Crutches

Some operations will require you to use crutches post-operatively. You will be made aware of this before your surgery. If you think you will have difficulty in using crutches please let one of the team know, so that alternative walking aids can be arranged.

Death

This is extremely rare and most unlikely to happen, but as with all surgical procedures it remains a remote possibility e.g. as a result of an anaphylactic shock (severe allergic reaction) to an injection. The podiatric surgery team is trained, and the department equipped, to deal with such an emergency.

Delayed healing

Although you will be given an estimate as to how long the recovery process will take; delays in healing of the soft tissue or bone can occur. Generally, the soft tissue will heal although bone sometimes does not heal properly (see non union). You should remember that the information you have been given is a guide and you should allow for this when planning your recovery.

Deterioration of symptoms

The vast majority of patients undergoing foot surgery have less discomfort following their operation. Occasionally however, some patients have no improvement of symptoms and more rarely, some patients have more discomfort. It is for this reason that foot surgery for cosmetic reasons is not recommended. It is always possible to have a straighter toe or foot which hurts more.

Driving

You should not drive after your foot surgery, until advised by the Podiatric Surgeon or one of the team, as you may damage the operation site and to do so may invalidate your insurance and be an offence under the Road Traffic Act.

Failure of the operation

On average at least 80-90 patients out of 100 are pleased with result of their surgery. The number would depend on the type / complexity of the operation and health of the patient. The important point to note is that despite the very best efforts of the team we cannot guarantee outcomes. This is true of any patient undergoing any surgery. In some operations (e.g. bunion surgery) full correction may not be achieved although symptoms are usually improved.

Fixation problems

Any metalwork (pins / screw / plate / implant) used, is normally left in place. In about 10% of patients metalwork needs to be removed (if it works loose or cause irritation to the surrounding tissues. Once bone is healed, metalwork is not required but unless it causes a problem we leave it in place.

General Anaesthetic

General anaesthetic means you will be ‘asleep’ during your operation. There is a slightly higher risk of medical complications for patients receiving general anaesthetic compared to local anaesthetic. Some patients may feel nauseous ‘sick’ when they wake up from the general anaesthetic.

THIS INFORMATION IS INTENDED AS A GUIDE ONLY

It is designed to be considered in conjunction with specific information concerning your particular operation

Infection

Infection rates are estimated at less than 2% of all surgeries. If infection does develop it is generally superficial, usually treated with antibiotics and resolves very easily. However, although uncommon, infection can get deeper into the bone which is a much more serious complication. This may require a period of hospitalisation, possibly further surgery and a longer course of antibiotics. In rare instances, the infected bone may need removing which could affect your foot permanently. Any infection has the potential to be fatal and foot surgery is no exception. Fortunately, this is extremely rare.

Joint stiffness

Surgery near joints e.g. bunion surgery can lead to some joint stiffness. The immobilisation following surgery and healing of the deep tissues near to the joint may be the underlying cause. We will usually give you some gentle exercises to minimise this risk.

Keloids

Keloids are large unsightly scars which are raised up and may extend beyond the margins of the original wound. This is a rare type of scar. Patients may have a history of poor scar formation / keloids, although having had no previous history of keloids does not mean you will not develop one. It is recognised that certain ethnic backgrounds may predispose to (increase the chance of developing) this type of scar e.g. Afro-Caribbean. In patients who have a history of ‘keloids’ we need to weigh-up very carefully the real risk of ‘keloid’ developing after surgery. There are some things we can try to minimise the risk but they are not 100% effective. These measures include:

- Use of corticosteroid injections prior to surgery
- Careful incision planning
- Use of zinc creams postoperatively

Local Anaesthetic

Local anaesthetics are medicines which when injected near nerves stop pain. Most of us have experienced a local anaesthetic at the dentist. Foot surgery is readily performed under local anaesthetic and involves injections placed either around the ankle or around the knee. After the injection has had time to work, which can take several minutes, you will not normally be able to feel pain in the area affected. A small number of patients do require additional local anaesthetic during surgery. In rare circumstances some patients do not respond to local anaesthetic injections and we cannot proceed with surgery (< 1 case per 100). In these situations surgery may have to be postponed. Whilst local anaesthetic drugs are considered very safe there are some potential side-effects such as:

- Allergic reaction to the anaesthetic (< 1 person per 10,000)
- Toxic reaction to the drug (< 1 person per 2000 across all injections)
- Irritation or damage to the nerves at the injections sites
- Bruising around the injection site (1 person in 20)

Loss of sensation

It is possible for you to lose some sensation around or away from the surgical site after surgery. Great care is taken to avoid damage to nerves and it is rare for a major nerve to be damaged. If nerve damage does occur it is more often the tiny nerves to the skin which may leave an area of skin with reduced or altered sensation. This loss of sensation can sometimes recover with time but this is not always the case.

Loss of tissue / part of foot / limb

This would be a very rare complication for the vast majority of patients having foot surgery. However it is possible for injury to blood vessels and or serious infection to lead to loss of tissue which can involve some or all of the foot. The risk of this for healthy patients would be much less than 1%.

THIS INFORMATION IS INTENDED AS A GUIDE ONLY

It is designed to be considered in conjunction with specific information concerning your particular operation

Metatarsalgia

This refers to discomfort under the ball of the foot (metatarsal heads). Surgery to the foot may alter the pressure under the forefoot and increase discomfort here. Sometimes this is treated with shoe inserts and rarely by further surgery.

Non-Union

This term refers to the situation where bone fails to fuse (join) together as planned. The percentage risk varies greatly depending on the bone or joint in question e.g. Bunion surgery risk is less than 1% and major joint fusion such as talo-navicular joint (big joint near ankle) risk is higher around 10% - 20%
If this situation occurs we may:

- Allow more time for the bones to knit together
- Re-operate on the bone / joint using a graft of bone from another part of your foot / body
- Do nothing if there are no symptoms as a result

Numbers of Operations Performed

In most cases your treating Consultant or Podiatric Surgeon will have performed a large number of the same operation for which you have been scheduled. In some circumstances however the consultant may have only performed a very small number of such operations. This might be for example if your operation is not something that is commonly required to be performed. Also sometimes the combinations of different operations necessary to fix your foot may not be commonly required.

On-call availability

On-call arrangements vary depending upon where you live. Some Podiatric Surgery services provide a specific on-call service, alternatively this may be provided via your GP out-of-hours on-call service, A&E or via another arrangement. This information will be provided when you are discharged from the hospital / day-surgery unit.

Oral Contraceptive medication

Certain (oestrogen containing) oral contraceptive medications are associated with a slight increased risk of thrombosis. You may wish to discuss the implications of this with one of the Podiatric Surgery team or your own GP. Stopping oral contraceptives means you are at risk of pregnancy unless alternative contraceptive measures are taken. Continuing with some types of oral contraceptives may mean you are at an increased risk of a blood clot.

Pain

It is difficult to predict how much pain you will suffer after the operation as this is variable between patients. Generally, the first night is the worse night but advances with local anaesthetics and pain medication means that this can be managed if not avoided.

Pain Medication

You may be advised to take medicines after your operation to help control postoperative pain.

Postoperative pain

Most people find they have mild to moderate pain after their surgery. The level of pain varies between individuals according to pain threshold and what medications patients are able to take. You will be provided with instructions on what to do should you have any problems immediately after your surgery.

Recurrence of the original problem

It is possible for a problem to recur postoperatively for a number of operations. For example a bunion corrected and looking perfect at 2 weeks postoperatively may deteriorate overtime. Recurrence varies greatly for individual operations.

THIS INFORMATION IS INTENDED AS A GUIDE ONLY

It is designed to be considered in conjunction with specific information concerning your particular operation

Rest

You are strongly advised to rest after your surgery. Resting with your foot elevated with an ice pack over the ankle will greatly reduce pain and swelling.

Scars

All surgery will create scars. Great care is taken to minimise the scar you will have by carefully placing the incision and taking great care during your operation. Some people will have a discomfort, nerve entrapment or thickening of the scar. These are uncommon, generally short term and, if they occur, they are often related to postoperative infection or to a predisposition to problem scars (see also Keloids). The vast majority of patients have minimal cosmetic defect from their scars. If you have any concerns please speak with one of the team for further advice.

Second Opinion

We are very happy to arrange for you to have a second opinion regarding your diagnosis or proposed treatment at any stage. Please notify one of our staff if you feel you would like a second opinion. Seeking a second opinion does not mean we will not treat you if you decide you wish to continue to receive care from within our department.

Secondary Arthritis

Some operations, generally joint fusions, lead to an increased risk of arthritis developing in adjacent joints. This is termed “*secondary arthritis*”. Fusion operations are generally performed to deal with pain from arthritic joints beyond salvage or because of instability in the foot. This can place more stress on adjacent joints increasing the risk of secondary arthritis in these joints.

Shoe difficulty

Although every effort is made to give as good a result as possible, you may still have difficulty with shoes and, in rare cases have less shoe choice after the operation.

Smoking

Smoking has the following adverse effects in relation to surgery:

- Delays wound healing
- Is associated with failure of bones to fuse [‘knit together’]. Risk increase 2.7 times more compared with a non-smoker
- Is associated with increased risk of thrombosis

Swelling

Swelling is always present after surgery as it is part of the normal healing process. Swelling may be minimised by following the post-operative instructions issued to you by the team. Some patients experience prolonged swelling of the foot after surgery (5-10%). Smaller operations usually give rise to minimal swelling whereas larger operations may be associated with greater swelling for longer duration. In a few cases, swelling may be present long term but this is usually painless although can affect shoe fit.

Success / Failure Rates

Many patients wish to know the overall success rates of individual operations. Sometimes we are able to provide this information based on our own audit data. Sometimes however, where it is a more unusual operation, statistics may be based on those from the literature. All operations carry risks and patients must accept this fact before electing to proceed with surgery.

THIS INFORMATION IS INTENDED AS A GUIDE ONLY

It is designed to be considered in conjunction with specific information concerning your particular operation

Thrombosis

Thrombosis has been the subject of much discussion recently in relation to flying. A thrombosis is a clot which most often forms in veins of the lower legs e.g. 'Deep Vein Thrombosis'. Risk factors for thrombosis are:

- Previous history of thrombosis
- Individual predisposition
- Smoking
- Certain drugs (e.g. some types of oral contraceptives)
- Damage to vessels and tissue (e.g. surgery)
- Immobilisation (including recent air travel i.e. within in previous 2 weeks)

The post surgery thrombosis risk has been calculated at 0.5%, however the more risk factors you possess the greater the chance of a clot forming. In patients with a history of clots, medicines to help reduce the risk of the further clots following your surgery will be arranged. In other circumstances you should reduce what risk factors you can by:

- Stop smoking at least 4 weeks prior to you operation and until you are recovered
- Avoid medicines which predispose to clots e.g. oral contraceptives (in discussion with your G.P, who will advise you of alternative measures of contraception)
- Follow the exercise regime as recommended by your podiatric surgeon or the team

Trainees

Many Podiatric Surgery services are approved clinical training centres for the Faculty of Surgery, College of Podiatrists. If you do not wish to have a trainee participate in your care; please notify a member of staff. Making this request will NOT in anyway disadvantage you.

Uneventful Recovery

The vast majority of patients who undergo foot surgery under the care of a Podiatric surgeon have an uneventful experience and are pleased with the result of their surgery. This booklet is designed to make sure that patients are fully aware of the potential for complications despite their low frequency. Some patients may feel more anxious being provided with this information but it is hoped that the number is low.

It is hoped that this booklet has answered all of your questions about your proposed operation. If you feel there is anything else you need to know, please do not hesitate to contact the podiatric surgery team.

PLEASE CONTACT ME IF YOU NEED ANY FURTHER INFORMATION OR ADVICE

THIS INFORMATION IS INTENDED AS A GUIDE ONLY

It is designed to be considered in conjunction with specific information concerning your particular operation